INTEDE: interactome of drug-metabolizing enzymes

Jiayi Yin^{1,†}, Fengcheng Li^{1,†}, Ying Zhou^{2,†}, Minjie Mou^{1,†}, Yinjing Lu¹, Kangli Chen¹, Jia Xue¹, Yongchao Luo¹, Jianbo Fu¹, Xu He¹, Jianqing Gao^{1,3}, Su Zeng^{1,3,*}, Lushan Yu^{1,*} and Feng Zhu [©]^{1,3,*}

¹College of Pharmaceutical Sciences, Zhejiang University, Hangzhou 310058, China, ²The First Affiliated Hospital, Zhejiang University, Hangzhou 310000, China and ³Innovation Institute for Artificial Intelligence in Medicine of Zhejiang University, Hangzhou 310018, China

Received July 20, 2020; Revised August 19, 2020; Editorial Decision August 25, 2020; Accepted September 22, 2020

ABSTRACT

Drug-metabolizing enzymes (DMEs) are critical determinant of drug safety and efficacy, and the interactome of DMEs has attracted extensive attention. There are 3 major interaction types in an interactome: microbiome-DME interaction (MICBIO), xenobiotics-DME interaction (XEOTIC) and host protein-DME interaction (HOSPPI). The interaction data of each type are essential for drug metabolism, and the collective consideration of multiple types has implication for the future practice of precision medicine. However, no database was designed to systematically provide the data of all types of DME interactions. Here, a database of the Interactome of Drug-Metabolizing Enzymes (INTEDE) was therefore constructed to offer these interaction data. First, 1047 unique DMEs (448 host and 599 microbial) were confirmed, for the first time, using their metabolizing drugs. Second, for these newly confirmed DMEs, all types of their interactions (3359 MICBIOs between 225 microbial species and 185 DMEs; 47 778 XEOTICs between 4150 xenobiotics and 501 DMEs; 7849 HOSPPIs between 565 human proteins and 566 DMEs) were comprehensively collected and then provided, which enabled the crosstalk analysis among multiple types. Because of the huge amount of accumulated data. the INTEDE made it possible to generalize key features for revealing disease etiology and optimizing clinical treatment. INTEDE is freely accessible at: https://idrblab.org/intede/

GRAPHICAL ABSTRACT



INTRODUCTION

Drug-metabolizing enzymes (DMEs) transform the parent drug to a metabolite with substantially different physicochemical and pharmacological properties, and are critical determinants of drug safety and efficacy (1,2). The interactome of DMEs has therefore attracted considerable attention (3-12). There are three major types of interactions: (i) microbiome-DME interactions (MICBIOs) determining the dynamic nature (5) and interpersonal variability (6) of drug metabolism; (ii) host protein–DME interactions (HOSPPIs) essential for predicting in vivo efficacy or clearance based on *in vitro* result (7) and revealing the mechanism underlying drug resistance (8) or adverse drug reaction (9); (iii) xenobiotics–DME interactions (XEOTICs) that are the key factor of metabolism-based drug-drug interaction (5) and a constant inspiration of clinical treatment optimization (10). Because the metabolism of drugs is collec-

*To whom correspondence should be addressed. Tel: +86 189 8946 6518; Fax: +86 571 8820 8444; Email: zhufeng@zju.edu.cn

(http://creativecommons.org/licenses/by-nc/4.0/), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Correspondence may also be addressed to Prof. Su Zeng. Email: zengsu@zju.edu.cn

Correspondence may also be addressed to Prof. Lushan Yu. Email: yuls@zju.edu.cn

[†]The authors wish it to be known that, in their opinion, the first four authors should be regarded as Joint First Authors.

[©] The Author(s) 2020. Published by Oxford University Press on behalf of Nucleic Acids Research.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License



Figure 1. INTEDE statistics of (i) the DMEs, and their corresponding drug(s) and tissue/disease specific protein abundances (*orange*) and (ii) three types of DME interactions: MICBIOs (*green*), HOSPPIs (*blue*) and XEOTICs (*red*). MICBIO determines the evolving and dynamic nature of metabolizing capacity and the interpersonal variability of drug metabolism; HOSPPI is essential for predicting *in vivo* efficacy/clearance based on the *in vitro* data, and revealing the mechanisms underlying drug resistance and toxicity; XEOTIC is one of the leading factors of the metabolism-based drug-drug interaction and the constant inspiration of clinical treatment optimization.

tively determined by multiple types of interactions, the accumulation of DME interactome data may give vital insight into the prediction of clinical consequences (7) and will have implication for the future practice of precision medicine (5).

Moreover, because of its dynamic evolvements (5), wide distribution (13), and rich repository of enzymes (14), the microbiome and its DMEs are found to be not only as critical as human DMEs in drug metabolism (10), but also key in the study of individual or tissue-specific metabolism of drugs (13,14) and the discovery of novel therapeutics targeting microbial protein (10). As a result, there was an explosive growth of interest in studying the interactome of microbial DME (15–19). Particularly, the MICBIO between actinobacteria *Eggerthella lenta* and microbial DME tyrosine decarboxylases was found to collectively regulate the metabolism of levodopa (10); the HOSPPI between host UDP-glucuronosyltransferases and microbial DME beta-glucuronidases could lead to an adverse drug reaction of irinotecan (20); the XEOTIC between antibiotic ciprofloxacin and bacterial DME cytidine deaminase was discovered to abrogate the resistance of gencitabine (21). Thus, it is important to collect these interactome data, and collectively analyze the multiple types of interactions for not only human but also microbial DMEs (7,10,15).

Nowadays, various databases have been developed to provide DME-related data, the majority of which are freely accessible and remain active. Some of these databases (such as DrugBank (22), PDB (23), PharmGKB (24), Protein Atlas (25), TTD (26) and UniProt (27)) provide the enzyme or DME information as part of a broader collection of biological/pharmacological data, and some others (including BRENDA (28), KEGG (29), MetaCyc (30), Reactome (31), SMPDB (32) and VMH (33)) contain the general enzymatic classifications or metabolic reaction/pathway data for a comprehensive set of enzymes. Since all these databases do not include DME interactome data, some databases have been designed to provide the interactions between ~2800 xenobiotics and ~190 DMEs (including CTD (34), T3DB (35) and Transformer (36)). However, no database has been designed to describe the key role of microbial DME in drug metabolism, and the interaction data of neither microbiome (MICBIO) nor host protein (HOSPPI), that alter drug metabolism by affecting human/microbial DME, are provided in any available knowledge base. As the crosstalk among interaction types is crucial for drug metabolism (7,10,15), it is critical to construct a database that comprehensively describes all three types of DME interactions.

Here, a newly developed database, Interactome of Drug-Metabolizing Enzyme (INTEDE) was therefore introduced. First, a comprehensive literature review of all (>1900) drugs approved by U.S. FDA and >3000 drugs in clinical trial or preclinical research was performed. Different from the small amount of DMEs (\sim 30) that were officially described in the U.S. FDA guidance (37,38), 553 DMEs (241 and 312 from human and microbiome, respectively) were confirmed in INTEDE to metabolize approved drugs, 421 DMEs (188 from human and 233 from microbiome) were to metabolize the drug in clinical or preclinical test, and 494 DMEs (208 from human and 286 from microbiome) were to metabolize the pharmaceutically investigative agents. Second, for all these newly confirmed DMEs, INTEDE comprehensively provided all three types of their interactions (MICBIOs, HOSPPIs and XEOTICs, as shown in Figure 1), which allowed the crosstalk among multiple interaction types (7,10,15) and could thus facilitate the study of disease etiology and the optimization of clinical treatment. All in all, since such crosstalk is key for drug metabolism, the INTEDE is expected to have implications for the future practice of precision medicine.

FACTUAL CONTENT AND DATA RETRIEVAL

Confirmation and collection of human/microbial DMEs

The DMEs collected in INTEDE were first confirmed by the drugs of clinical importance (either approved or clinical/preclinical/investigative). Particularly, a comprehensive literature review of all drugs approved by FDA (1921 in total, collected from the U.S. FDA official site), 2958 drugs in clinical trial (data from the ClinicalTrial.gov (39) and TTD (40)) and 10 213 pharmaceutically investigative agents (obtained from the TTD (40)), was performed to confirm their corresponding DMEs by searching the PubMed (41) using such keyword combinations as "Drug Name' + drug metabolizing enzyme', "Drug Name' + metabolism', "Drug Name' + enzyme', "Drug Name' + drug metabolism' and "Drug Name' + metabolized'. Moreover, to facilitate the discovery of microbial DMEs, >2000 microbial species (from bacteria, fungi, metamonada, amoebozoa and archaea (3,42,43) colonizing throughout the human body (such as bladder, blood, eye, gut, lung, oral cavity, skin, stomach, urethra and vagina (44-46)) were collected, and the microbial DMEs were then discovered by the literature review of PubMed (41) using such keyword combinations as "Species Name' + 'Drug Name'', "Species Name' + drug metabolism', "Species Name' + drug metabolized' and "Species Name' + metabolizing enzyme'.

As a result, 553 DMEs (241 host and 312 microbial) were identified to metabolize 924 approved drugs, 421 DMEs

(188 host and 233 microbial) were to metabolize 560 drugs in preclinical and clinical, and 494 DMEs (208 host and 286 microbial) were to metabolize 405 pharmaceutically investigative agents. In total, there were 1047 unique DMEs in INTEDE (448 from human and 599 from microbes). On one hand, 448 human DMEs were grouped to diverse enzymatic families (43 families in total defined by the second level of the Enzyme Commission (EC) nomenclature (28)), and the Top-5 popular DME families were: Paired Donor Oxidoreductase (EC: 1.14, 67 DMEs), CH-OH Donor Oxidoreductase (EC: 1.1, 47 DMEs), Ester Bond Hydrolase (EC: 3.1, 41 DMEs), Glycosyltransferase (EC: 2.4, 34 DMEs) and Kinase (EC:2.7, 32 DMEs). On the other hand, 599 microbial DMEs also belonged to diverse enzymatic families (29 families in total defined by the second level of EC nomenclature), and the Top-5 popular DME families were: Carbon-Nitrogen Hydrolase (EC: 3.5, 99 DMEs), Glycosylase (EC: 3.2, 91 DMEs), Paired Donor Oxidoreductase (EC: 1.14, 33 DMEs), CH-NH Donor Oxidoreductase (EC: 1.5, 31 DMEs), and Acyltransferase (EC: 2.3, 20 DMEs). Experimentally assessed kinetic parameters (Km) and catalytic efficiencies (Kcat/Km) between 235 drugs and 190 human/microbial DMEs were also offered. As shown in Figure 2, a full list of drugs metabolized by certain DME were displayed in INTEDE.

Tissue and disease specific protein sbundances of DMEs

Tissue and disease specific DME abundances are critical for the mitigation of drug toxic reaction (47) and alteration of drug pharmacokinetics (48). In particular, tissue specific DME abundances can not only maintain the functional homeostasis of drug, but also mitigate their tissue-dependent adverse reactions, such as cardiotoxicity, nephrotoxicity and neurotoxicity (47); the variations in DME abundances among different disease indications (such as infection, immunodeficiency and inflammation) can significantly affect hepatic/renal clearance, and therefore reduce drug efficacy or aggravate adverse drug reaction (48). Since these tissue/disease-dependent variations in DME abundances were expected to provide new therapeutic strategies (49-52), the relevant data should be accumulated and further analyzed to promote modern drug discovery.

Disease specific DME abundance data were collected as follows, the detailed procedure of which was illustrated in Supplementary Method S1. First, 5304 series records from the same microarray platform (Affymetrix HGU133 Plus 2.0) were gathered from the Gene Expression Omnibus (53). Second, the collected records were sequentially processed by data normalization, transformation, integration, perfect match correction, quantile, robust multiarray average and median polish (54). Third, the differential expression analysis (55) was conducted by comparing the DME abundance among sample groups (defined in Supplementary Method S1). Fourth, the violin-plot describing the differential expression pattern of the studied DMEs among sample groups was drawn (shown in Supplementary Figure S1). Furthermore, tissue specific DME abundances data were collected using the following procedure, and the detailed process was elaborated in Supplementary Method S1.

Details	of Drug-Metab	olizing Enz	yme (DME)		
General Info	rmation of DME (ID: [DME0001)			
DME Name	Cytochrome P450 3A4 (CYP3A4), Homo sapiens				
EC Number	EC: 1.14.14.55 (Click to Show/Hide the Complete EC Tree)				
Lineage	Species: Homo sapiens (Click to Show/Hide the Complete Species Lineage)				
Interactome	 Interactions between Microbiome an Interactions between Xenobiotics an Interactions between Host Protein ar 		d DME (MICBIO) S Jump to Detailed In ad DME (XEOTIC) Jump to Detailed In and DME (HOSPPI) Jump to Detailed In		teractome Data teractome Data teractome Data
Click to Sh	ow/Hide the Molecular/Fur	ctional Data (Seque	nce/Structure/Pathway/Function	on) of This DME	
Full List of D	rug(s) Metabolized b	v This DME			
Drugs Approved by FDA			Click to Show/Hide the Full List of Drugs: 546 Drugs		
Drugs in Phase 4 Clinical Trial			Click to Show/Hide the Full List of Drugs: 81 Drugs		
Drugs in Phase 3 Clinical Trial			Click to Show/Hide the Full List of Drugs: 71 Drugs		
Drugs in Phase 2 Clinical Trial			Click to Show/Hide the Full List of Drugs: 50 Drugs		
Drugs in Phase 1 Clinical Trial			Click to Show/Hide the Full List of Drugs: 32 Drugs		
Preclinical/investigative Agents			Click to Show/Hide the Full List of Drugs: 41 Drugs		
Discontinued/withdrawn Drugs			Click to Show/Hide the Full List of Drugs: 22 Drugs		
Tissue/Disea	se-Specific Protein A	bundances of T	his DME		
Tissue-spe	cific Protein Abundances in	Healthy Individuals			Click to Show/Hide
ICD Disease Classification 01 Infectious/pa			rasitic disease		Click to Show/Hide
ICD Diseas	se Classification 02	Neoplasm			Click to Show/Hide
ICD Disease Classification 03 Blood/blog		Blood/blood-	-forming organ disease		Click to Show/Hide
ICD Disease Classification 04 Immune sys			em disease		Click to Show/Hide
ICD Diseas	se Classification 05	Endocrine/nu	tritional/metabolic disease		Click to Show/Hide
ICD Diseas	e Classification 06 Mental/behavioural/neurodevelopm			isorder	Click to Show/Hide
ICD Diseas	se Classification 08	Nervous syst	em disease		Click to Show/Hide
ICD Diseas	se Classification 09	Visual system	n disease		Click to Show/Hide
ICD Disease Classification 11 Circulatory set			ystem disease		Click to Show/Hide
ICD Disease Classification 12 Respiratory s			ystem disease		Click to Show/Hide
ICD Disease Classification 13 Digestive system			stem disease		Click to Show/Hide
ICD Disease Classification 14 Skin disease					Click to Show/Hide
ICD Disease Classification 15 Musculoskel			etal system/connective tissue disease		Click to Show/Hide
ICD Disease Classification 16 Genitourinar			system disease		Click to Show/Hide
ICD Disease Classification 19 Condition or			ginating in perinatal period		Click to Show/Hide
ICD Diseas	se Classification 20	Developmen	tal anomaly		Click to Show/Hide

Figure 2. A typical INTEDE page providing the details of DME, which included: (i) the general information (DME name, EC nomenclature, taxonomy lineage, hyperlink to detailed interactome and molecular or functional data); (ii) the full list of drugs metabolized by this DME (categorized based on clinical status: FDA approved, clinical, preclinical, investigative and discontinued. By clicking on each category, the detail information of each drug can be viewed); (iii) the tissue and disease specific protein abundance of the DME (the differential abundance profiles of 394 DMEs in 60 tissues and 106 diseases, and the expression profiles of 348 DMEs across 36 human tissues).

First, a benchmark dataset that contained the protein expression data across 36 human tissues (56) was collected. Second, the intensities of each DME were processed via data integration and scaling. Third, the expression bar plot of studied DME across 36 tissues was drawn (Supplementary Figure S2). Overall, the differential abundance profile of 394 DMEs in 20,663 patients and healthy individuals of 60 tissues and 106 diseases, and the expression profile of 348 DMEs across 36 human tissues were provided in and downloadable from the website. As shown in Figure 2, the tissue and disease specific DME abundances were described.

The interactome of human and microbial DMEs

INTEDE comprehensively provided all three types of DME interaction data (as shown in Figure 1) for those 1047 unique DMEs (448 from human and 599 from

microbe) confirmed in previous section. These interaction data included: (i) microbiome–DME interactions (MICBIOs); (ii) host protein–DME interactions (HOSP-PIs) and (iii) xenobiotics–DME interactions (XEOTICs).

Interactions between microbiome and DMEs. The microbial manipulation of drug metabolism by interacting with DME is characterized by (i) the vast diversity of involved microbial species (including various bacteria or fungi from dozens of taxonomic phyla (6,10,20,21,57–59)) and (ii) the wide distribution of these species throughout the human body (not only in gut, but also in skin, vagina, and many other sites (17,21,44,60–62)). Since the microbiome-DME interactions (MICBIOs) are key in determining the dynamic feature (5) and interpersonal variability (6) of drug metabolism, it is very essential to have the MICBIOs data for understanding the controls/effects of the circadian timing system (63) and individualized microbiota compositions (21,61) on the metabolism of certain medication (Figure 1).

Because the MICBIOs data were largely dispersed in literatures, PubMed database (41) was first searched to find the interactions between microbiome and human/microbial DMEs. In particular, the keyword combinations of "Microbial Species Name' + 'DME Name", "Species Name' + 'Drug Name" and "Species Name' + drug metabolism' were used in literature reviews, and the discovered publications were manually assessed for retrieving any MICBIOsrelated information. Moreover, due to the collective determination by multiple factors present in a drug delivery route (64), the metabolism of a drug should be assessed by simultaneously considering its DME(s) and the colonizing microbes, if both DME and microbe are involved in the same route (64). Therefore, additional literature reviews were then conducted using the keyword combination of "Microbial Species Name' + 'DME Name' + 'Drug Name". The studies that describe the DME and microbe that metabolized the same drug and present in the same delivery route were identified, and these publications were also manually evaluated for retrieving the MICBIOs-related data.

As a result, the collected data included the taxonomic lineage of each microbe (kingdom, phylum, class, order, family, genus and species), the human body site distributed by each microbe (bladder, blood, eye, gut, lung, oral cavity, skin, stomach, urethra and vagina), the list of drugs affected by the interaction between microbial species and DME, and the mechanism of how drug metabolism is influenced by the corresponding interactions. All in all, the latest version of INTEDE contained 3359 MICBIOs between 225 microbes and 185 human/microbial DMEs. These microbes came from 3 kingdoms, 19 phyla, 31 classes, 40 orders, 69 families and 107 genera as defined by the NCBI Taxonomy database (65), and colonized in 10 different human body sites (such as: bladder, blood, eye, gut, lung, oral cavity, skin, stomach, urethra and vagina). The MICBIOs data can be assessed and retrieved by various search strategies in both the Home page and the subpage entitled 'Microbiome-DME interaction' of INTEDE (as shown in Figure 3).

Interactions between host proteins and DMEs. Host protein-DME interactions (HOSPPIs) were frequently encountered in drug metabolism, and widely acknowledged to be critical determinant of drug safety and efficacy (7-9,66-69). A variety of HOSPPIs were of great interest in recent metabolic study, which included (i) oligomerizations that are essential for predicting in vivo drug efficacy or clearance based on the in vitro results (7), (ii) transcription factor regulations that are critical for revealing the mechanisms underlying drug resistance (8) and metabolic variation (70,71), (iii) epigenetic regulations (like DNA methylations, histone modifications and non-coding RNA regulations) that lead to inter-individual variability in drug responses and adverse reactions, and highlight the implications for personalized medicine (9,72,73). Furthermore, the effects of HOSPPIs on altering the drug metabolisms varied between healthy and pathological condition, and differed among disease indications (74). It was therefore essential to collect HOSPPIs to facilitate the prediction of clinical consequence (Figure 1).

The HOSPPIs data was identified by literature search using different keyword combinations. For example, when searching for the oligomerization of a DME, "DME Name" + oligomer', "DME Name' + oligomerization', "DME Name' + dimer', "DME Name' + trimer', "DME Name' + dimerization', "DME Name' + trimerization' and "DME Name' + protein-protein interaction' were used; when discovering the transcription factor regulations on a DME, 'transcription factor + 'DME Name" and 'transcriptional regulation + 'DME Name" were reviewed; when finding the DNA methylations of a DME, "DME Name' + methylation', "DME Name' + epigenetics' and "DME Name' + methylate' were searched. Moreover, the additional DNA methylation data for all DMEs in INTEDE were further collected using the following processes. First, 1377 series records of a very popular microarray platform (Illumina HumanMethylation450 BeadChip) were collected from the Gene Expression Omnibus (53), and 136 series records with both patients and healthy individuals data were selected, which contained 16 256 samples of 86 disease indications. Second, all these collected records were sequentially processed by quality control, normalization and batch effect correction (75-78), and differential methylations were discovered by calculating both P-values and delta-beta values (79,80). According to the definition in previous reports, the threshold of *P*-values was set to 0.05 (*P*-value < 0.05 was considered to be statistically significant (79)), and the cutoff of delta-beta values was set to 0.2 and 0.3 (the absolute value of *delta-beta* > 0.3 was considered as 'significant methylation' (79), and $0.3 \ge |delta-beta| > 0.2$ was reported as 'moderate methylation' (80)). Third, the above differential methylations were discovered among four different sample groups, which included: (i) DME methylation in the normal tissue adjacent to the disease tissue of patients (blue), (ii) DME methylation in the disease tissue of patients (red), (iii) DME methylation in the normal tissue of healthy individual (green) and (iv) DME methylation in the tissue other than the disease tissue of patients (yellow). Fourth, the methylation plot across samples was drawn using ggplot2 in R environment, and the violin plots of methylation variation between two studied groups was generated using ggplot2 and ggbeeswarm in R. All plots can be viewed online (Supplementary Figure S3) and downloaded from the website.

Consequently, the HOSPPI information included the interaction types (oligomerization, transcription factor regulation, DNA methylation and drug co-metabolism), the studied diseases (95 diseases standardized in a latest version of International Classification of Diseases (81), ICD-11), the targeted cell lines (289 cell lines, such as A549, Caco-2, HEK293, HepG2 and MCF-7), the levels of DNA methylation (significant or moderate) together with the corresponding *P*-value and *delta-beta* value, the drugs which metabolism was affected by studied HOSPPIs, the detailed mechanism of a HOSPPI on drug metabolism, and the violin/bar plots illustrating the differential methylation between different sample groups. All in all, INTEDE contained 7849 HOSPPIs between 565 host proteins and 566 DMEs: oligomerization (homo-/hetero-), transcription factor regulation (activation/repression), DNA methylation (hypo-/hyper-), drug co-metabolism (direct/indirect). HOSPPIs data can be assessed in INTEDE (Figure 4).



Figure 3. A typical INTEDE page offering the data of microbiome-DME interaction (MICBIO). The INTEDE contained 3359 MICBIOs between 225 microbes and 185 DMEs. These microbes came from 3 kingdoms, 19 phyla, 69 families and 107 genera, and colonized in 10 different host body sites (bladder, blood, eye, gut, lung, oral cavity, skin, stomach, urethra and vagina).

Interactions between xenobiotics and DMEs. The activity or expression of DMEs could be inhibited or induced by xenobiotics, which, in turn, affected the hepatic clearance of drug and led to the reduced efficacy and therapeutic failure (82–85). These xenobiotics included: (i) pharmaceutical agents (FDA approved, clinical, preclinical, patented and investigative), (ii) health toxicants (biotoxin, carcinogen, environmental pollutant, mycotoxin, and neurotoxin), (iii) natural substances (biological extract, natural product/mixture and traditional medicine), (iv) agricultural chemicals (fungicides, herbicides and insecticide), (v) additive agents (food additives and cosmetic additives) and (vi) proteins/peptides. Since all these xenobiotics are one of the leading factors of the metabolism-based drug-drug interactions (5) and the constant inspiration of clinical treatment optimizations (10), it was essential to collect those xenobiotics-DME interactions (XEOTICs), and clarify the effects of an XEOTIC on modulating the activity or expression of human/microbial DMEs.

Therefore, the XEOTIC data were systematically searched by literature review in PubMed based on the keyword combinations between 'DME Name' and 'biotoxin', 'environmental pollutant', 'extract', 'fungicides', 'inducer', 'inhibitor', 'insecticides', 'nature product' and 'pesticides'. Those discovered literatures were evaluated manually to identify the xenobiotics affecting DMEs. The collected data included the name of xenobiotics, the modulation types (inducer and inhibitor), the xenobiotic classifications (additive, agricultural chemical, health toxicant, natural substance, and pharmaceutical agent), the modulation activity (measured by MIC/IC50/Ki values), and the affected cell system (choriocarcinoma, hamster ovary, and high five). All in all, the INTEDE provided 47,778 XEOTICs between 4150 xenobiotics and 501 human or microbial DMEs. These miscellaneous xenobiotics cover all the xenobiotic types described in the previous paragraph (from (i) pharmaceutical agents to (vi) proteins/peptides).



Figure 4. A typical INTEDE page offering the data of host protein-DME interactions (HOSPPIs). The INTEDE contained 7849 HOSPPIs between 565 host proteins and 566 DMEs. Various types of HOSPPI were in the INTEDE, which included: oligomerization (homo-/hetero-), transcription factor regulation (activation/repression), and DNA methylation (hypo-/hyper-).

All XEOTICs can be assessed in the subpage of INTEDE entitled 'Xenobiotics-DME Interaction' (Figure 5).

Data standardization, access and retrieval

To make the access and analysis of INTEDE data convenient for all users, the collected raw data were carefully cleaned up and then were systematically standardized, which included the disease standardization, EC classification, unit unification, structure availability and crosslink to various reference databases. Detailed description on the way to clean and standardize data was provided in Supplementary Method S2. Furthermore, INTEDE has been smoothly running for months, and tested from different sites around the world. All data could be viewed, accessed, and downloaded (Supplementary Figure S4). Currently, the INTEDE is freely assessed without login requirement by all users at: https://idrblab.org/intede/.

INTERACTION CROSSTALK AND PERSPECTIVES

Although the data of each type of DME interaction in INTEDE were essential for current clinical studies, the

crosstalk among different interaction types had emerged to be increasingly important due to the extremely complex mechanisms underlying drug metabolism (10,20,57,86,87). Taking the primary medication for treating *Parkinson's disease*—levodopa—as an example, it must enter the brain and be converted there by the human DME aromatic amino acid decarboxylase (AADC) to its active form-dopamine. However, the delivery of levodopa to blood-brain barrier is greatly restricted by its intestinal metabolisms of both human and gut microbiome (10). Particularly, the host intestinal AADC and gut microbes Enterococcus fae*calis* and *Eggerthella lenta* can convert levodopa into a different chemical before it reaches the brain, which causes unwanted side effects (10). Therefore, levodopa should be co-administrated with other drug by collectively considering multiple DME interactions: (i) the MICBIO between the above microbes and brain DME AADC; (ii) the XEOTIC between xenobiotics carbidopa (inhibitor of intestinal AADC) and AADC (10). In other words, the crosstalk between the MICBIO and XEOTIC of human DME AADC inspired a new strategy that combines AFMT (a microbe inhibitor) and carbidopa (a DME inhibitor)



Figure 5. A typical INTEDE page offering the data of xenobiotics-DME interaction (XEOTIC). The INTEDE provided 47 778 XEOTICs between 4150 xenobiotics and 501 DMEs, which were diverse. Particularly, the types of xenobiotics included (i) pharmaceutical agents (FDA approved, clinical, preclinical and investigative), (ii) health toxicants (biotoxin, carcinogen, environmental pollutant, mycotoxin and neurotoxin), (iii) natural substances (biological extract, natural product and traditional medicine), (iv) agricultural chemicals (fungicides, herbicides and insecticide), (v) additive agents (food additive and cosmetic additive) and (vi) proteins /peptides.

with levodopa, for effectively overcoming the serious side effects of anti-*Parkinson* drug (10).

Similarly, the HOSPPI between host protein carboxylesterase and DME CYP3A4 was found as the leading cause of drug resistance, which could therefore be reversed by the XEOTIC between xenobiotics vitamin D and DME CYP3A4 (87); the HOSPPI between host protein UGT1A1 and microbial DME beta-glucuronidase was reported to cause diarrhea, which needed be relieved by the XEOTIC between xenobiotic amoxapine and the microbial DME beta-glucuronidase (20,57). Thus, the multiple types of DME interaction in INTEDE might keep inspiring new strategies for dealing with drug resistances and side effects.

To make the crosstalk analysis of each DME possible, the data of multiple interaction types were systematically collected. As a result, 453 (43.3%) out of 1047 DMEs were described in INTEDE with multiple interaction types. Moreover, the number of interaction data for each type was

from thousands to tens of thousands, which made it highly possible to identify the differential features or generalize the common characteristics from these DME-related big data. Meanwhile, because of the lack of multiple interaction types for the remaining 56.7% DMEs, it is essential to continue drug metabolism studies for a further extension of our knowledge on DME interactome.

In summary, INTEDE is unique in: (i) providing the largest number of DMEs that are manually curated and systematically confirmed based on their metabolizing drug(s), (ii) covering the novel DME data of diverse microbes (parasitizing in different sites of human body) together with their metabolizing drug(s), (iii) describing the comprehensive interactome data from three perspectives for both human and microbial DMEs and (iv) enabling the crosstalk analysis among various types of DME interaction. With the extensive efforts made on describing DME interactome (3–10) and revealing the crosstalk among interaction types (10,20,57,86,87), those immense, connected and structural-

ized data provided in INTEDE are expected to have implications for the future practice of clinical treatment optimization (88–92) and personalized medicine (93–96).

SUPPLEMENTARY DATA

Supplementary Data are available at NAR Online.

FUNDING

National Key R&D Program of China [2017YFC0908600, 2018YFC0910500]; National Natural Science Foundation of China [U1909208, 81872798]; Leading Talent of the 'Ten Thousand Plan' – National High-Level Talents Special Support Plan, Key R&D Program of Zhejiang Province [2020C03010]; Fundamental Research Funds for the Central Universities [2018QNA7023, 10611CD-JXZ238826, 2018CDQYSG0007, CDJZR14468801]; Information Technology Center, Zhejiang University; China Knowledge Centre for Engineering Sciences and Technology [No. CKCEST-2019-1-12]. Funding for open access charge: National Natural Science Foundation of China [81872798].

Conflict of interest statement. None declared.

REFERENCES

- Kirchmair, J., Goller, A.H., Lang, D., Kunze, J., Testa, B., Wilson, I.D., Glen, R.C. and Schneider, G. (2015) Predicting drug metabolism: experiment and/or computation? *Nat. Rev. Drug Discov.*, 14, 387–404.
- 2. Hitchings, R. and Kelly, L. (2019) Drug metabolism as a community effort. *Cell Metab.*, **30**, 235–237.
- Almeida, A., Mitchell, A.L., Boland, M., Forster, S.C., Gloor, G.B., Tarkowska, A., Lawley, T.D. and Finn, R.D. (2019) A new genomic blueprint of the human gut microbiota. *Nature*, 568, 499–504.
- Scott, T.A., Quintaneiro, L.M., Norvaisas, P., Lui, P.P., Wilson, M.P., Leung, K.Y., Herrera-Dominguez, L., Sudiwala, S., Pessia, A., Clayton, P.T. *et al.* (2017) Host-microbe co-metabolism dictates cancer drug efficacy in C. elegans. *Cell*, 169, 442–456.
- 5. Clarke, G., Sandhu, K.V., Griffin, B.T., Dinan, T.G., Cryan, J.F. and Hyland, N.P. (2019) Gut reactions: breaking down
- xenobiotic-microbiome interactions. *Pharmacol. Rev.*, **71**, 198–224.
 Zimmermann, M., Zimmermann-Kogadeeva, M., Wegmann, R. and Goodman, A.L. (2019) Mapping human microbiome drug
- metabolism by gut bacteria and their genes. *Nature*, 570, 462–467.
 7. Hu,D.G., Hulin,J.U., Nair,P.C., Haines,A.Z., McKinnon,R.A., Mackenzie,P.I. and Meech,R. (2019) The UGTome: the expanding diversity of UDP glycosyltransferases and its impact on small
- and the impact of ODP glycosyntansterases and its impact on small molecule metabolism. *Pharmacol. Ther.*, **204**, 107414.
 8. Xu,D., Huang,S., Wang,H. and Xie,W. (2018) Regulation of brain
- drug metabolizing enzymes and transporters by nuclear receptors. *Drug Metab. Rev.*, **50**, 407–414.
 9. Lauschke, V.M., Barragan, I. and Ingelman-Sundberg, M. (2018)
- Pharmacoepigenetics and toxicoepigenetics: novel mechanistic insights and therapeutic opportunities. *Annu. Rev. Pharmacol. Toxicol.*, **58**, 161–185.
- Maini Rekdal, V., Bess, E.N., Bisanz, J.E., Turnbaugh, P.J. and Balskus, E.P. (2019) Discovery and inhibition of an interspecies gut bacterial pathway for levodopa metabolism. *Science*, **364**, eaau6323.
- Kovalchuk, N., Zhang, Q.Y., Kelty, J., Van Winkle, L. and Ding, X. (2019) Toxicokinetic interaction between hepatic disposition and pulmonary bioactivation of inhaled naphthalene studied using Cyp2abfgs-null and CYP2A13/2F1-humanized mice with deficient hepatic cytochrome P450 activity. *Drug Metab. Dispos.*, 47, 1469–1478.
- Ma,J., Xia,Q., Fu,P.P. and Lin,G. (2018) Pyrrole-protein adducts a biomarker of pyrrolizidine alkaloid-induced hepatotoxicity. J. Food Drug. Anal., 26, 965–972.

- Wilkinson, E.M., Ilhan, Z.E. and Herbst-Kralovetz, M.M. (2018) Microbiota-drug interactions: impact on metabolism and efficacy of therapeutics. *Maturitas*, 112, 53–63.
- Doestzada, M., Vila, A.V., Zhernakova, A., Koonen, D.P.Y., Weersma, R.K., Touw, D.J., Kuipers, F., Wijmenga, C. and Fu, J. (2018) Pharmacomicrobiomics: a novel route towards personalized medicine? *Protein Cell*, 9, 432–445.
- Zimmermann, M., Zimmermann-Kogadeeva, M., Wegmann, R. and Goodman, A.L. (2019) Separating host and microbiome contributions to drug pharmacokinetics and toxicity. *Science*, 363, eaat9931.
- Eisenstein, M. (2020) Microbial matters: modelling the complex web of host-microbiome interactions. *Nature*, 581, 489–490.
- Lam,K.N., Alexander,M. and Turnbaugh,P.J. (2019) Precision medicine goes microscopic: engineering the microbiome to improve drug outcomes. *Cell Host Microbe*, 26, 22–34.
- Cani, P.D. (2018) Human gut microbiome: hopes, threats and promises. *Gut*, 67, 1716–1725.
- Savage,N. (2020) The complex relationship between drugs and the microbiome. *Nature*, 577, S10–S11.
- Collins,S.L. and Patterson,A.D. (2020) The gut microbiome: an orchestrator of xenobiotic metabolism. *Acta Pharm. Sin. B*, 10, 19–32.
- Geller, L.T., Barzily-Rokni, M., Danino, T., Jonas, O.H., Shental, N., Nejman, D., Gavert, N., Zwang, Y., Cooper, Z.A., Shee, K. *et al.* (2017) Potential role of intratumor bacteria in mediating tumor resistance to the chemotherapeutic drug gencitabine. *Science*, 357, 1156–1160.
- Wishart, D.S., Feunang, Y.D., Guo, A.C., Lo, E.J., Marcu, A., Grant, J.R., Sajed, T., Johnson, D., Li, C., Sayeeda, Z. et al. (2018) DrugBank 5.0: a major update to the DrugBank database for 2018. Nucleic Acids Res., 46, D1074–D1082.
- Burley,S.K., Berman,H.M., Bhikadiya,C., Bi,C., Chen,L., Di Costanzo,L., Christie,C., Dalenberg,K., Duarte,J.M., Dutta,S. *et al.* (2019) RCSB Protein Data Bank: biological macromolecular structures enabling research and education in fundamental biology, biomedicine, biotechnology and energy. *Nucleic Acids Res.*, 47, D464–D474.
- Barbarino, J.M., Whirl-Carrillo, M., Altman, R.B. and Klein, T.E. (2018) PharmGKB: a worldwide resource for pharmacogenomic information. *Wiley Interdiscip. Rev. Syst. Biol. Med.*, 10, e1417.
- Uhlen, M., Fagerberg, L., Hallstrom, B.M., Lindskog, C., Oksvold, P., Mardinoglu, A., Sivertsson, A., Kampf, C., Sjostedt, E., Asplund, A. *et al.* (2015) Tissue-based map of the human proteome. *Science*, 347, 1260419.
- 26. Wang, Y., Zhang, S., Li, F., Zhou, Y., Zhang, Y., Wang, Z., Zhang, R., Zhu, J., Ren, Y., Tan, Y. *et al.* (2020) Therapeutic target database 2020: enriched resource for facilitating research and early development of targeted therapeutics. *Nucleic Acids Res.*, **48**, D1031–D1041.
- UniProt,C. (2019) UniProt: a worldwide hub of protein knowledge. Nucleic Acids Res., 47, D506–D515.
- Jeske, L., Placzek, S., Schomburg, I., Chang, A. and Schomburg, D. (2019) BRENDA in 2019: a european ELIXIR core data resource. *Nucleic Acids Res.*, 47, D542–D549.
- Kanehisa, M., Sato, Y., Furumichi, M., Morishima, K. and Tanabe, M. (2019) New approach for understanding genome variations in KEGG. *Nucleic Acids Res.*, 47, D590–D595.
- Caspi, R., Billington, R., Keseler, I.M., Kothari, A., Krummenacker, M., Midford, P.E., Ong, W.K., Paley, S., Subhraveti, P. and Karp, P.D. (2020) The MetaCyc database of metabolic pathways and enzymes - a 2019 update. *Nucleic Acids Res.*, 48, D445–D453.
- Jassal, B., Matthews, L., Viteri, G., Gong, C., Lorente, P., Fabregat, A., Sidiropoulos, K., Cook, J., Gillespie, M., Haw, R. *et al.* (2020) The reactome pathway knowledgebase. *Nucleic Acids Res.*, 48, D498–D503.
- Jewison, T., Su, Y., Disfany, F.M., Liang, Y., Knox, C., Maciejewski, A., Poelzer, J., Huynh, J., Zhou, Y., Arndt, D. *et al.* (2014) SMPDB 2.0: big improvements to the Small Molecule Pathway Database. *Nucleic Acids Res.*, 42, D478–D484.
- 33. Noronha,A., Modamio,J., Jarosz,Y., Guerard,E., Sompairac,N., Preciat,G., Danielsdottir,A.D., Krecke,M., Merten,D., Haraldsdottir,H.S. *et al.* (2019) The virtual metabolic human database: integrating human and gut microbiome metabolism with nutrition and disease. *Nucleic Acids Res.*, 47, D614–D624.
- Davis, A.P., Grondin, C.J., Johnson, R.J., Sciaky, D., McMorran, R., Wiegers, J., Wiegers, T.C. and Mattingly, C.J. (2019) The comparative

toxicogenomics database: update 2019. Nucleic Acids Res., 47, D948–D954.

- Wishart, D., Arndt, D., Pon, A., Sajed, T., Guo, A.C., Djoumbou, Y., Knox, C., Wilson, M., Liang, Y., Grant, J. et al. (2015) T3DB: the toxic exposome database. *Nucleic Acids Res.*, 43, D928–D934.
- Hoffmann, M.F., Preissner, S.C., Nickel, J., Dunkel, M., Preissner, R. and Preissner, S. (2014) The Transformer database: biotransformation of xenobiotics. *Nucleic Acids Res.*, 42, D1113–D1117.
- 37. Taskar,K.S., Pilla Reddy,V., Burt,H., Posada,M.M., Varma,M., Zheng,M., Ullah,M., Emami Riedmaier,A., Umehara,K.I., Snoeys,J. *et al.* (2020) Physiologically-based pharmacokinetic models for evaluating membrane transporter mediated drug-drug interactions: current capabilities, case studies, future opportunities, and recommendations. *Clin. Pharmacol. Ther.*, **107**, 1082–1115.
- Iwatsubo,T. (2020) Evaluation of drug-drug interactions in drug metabolism: differences and harmonization in guidance/guidelines. *Drug Metab. Pharmacokinet.*, 35, 71–75.
- 39. Tse, T., Fain, K.M. and Zarin, D.A. (2018) How to avoid common problems when using ClinicalTrials.gov in research: 10 issues to consider. *BMJ*, **361**, k1452.
- 40. Li,Y.H., Yu,C.Y., Li,X.X., Zhang,P., Tang,J., Yang,Q., Fu,T., Zhang,X., Cui,X., Tu,G. *et al.* (2018) Therapeutic target database update 2018: enriched resource for facilitating bench-to-clinic research of targeted therapeutics. *Nucleic Acids Res.*, 46, D1121–D1127.
- Sayers, E. W., Beck, J., Brister, J.R., Bolton, E.E., Canese, K., Comeau, D.C., Funk, K., Ketter, A., Kim, S., Kimchi, A. *et al.* (2020) Database resources of the national center for biotechnology information. *Nucleic Acids Res.*, 48, D9–D16.
- 42. Hallen-Adams, H.E. and Suhr, M.J. (2017) Fungi in the healthy human gastrointestinal tract. *Virulence*, **8**, 352–358.
- Kwa, M., Plottel, C.S., Blaser, M.J. and Adams, S. (2016) The intestinal microbiome and estrogen receptor-positive female breast cancer. J. Natl. Cancer Inst., 108, djw029.
- 44. Nichols, R.G., Peters, J.M. and Patterson, A.D. (2019) Interplay between the host, the human microbiome, and drug metabolism. *Hum. Genomics*, **13**, 27.
- Ozkan, J., Willcox, M., Wemheuer, B., Wilcsek, G., Coroneo, M. and Thomas, T. (2019) Biogeography of the human ocular microbiota. *Ocul. Surf.*, 17, 111–118.
- Simon,C. (2018) Introduction: do microbes in the female reproductive function matter? *Fertil. Steril.*, 110, 325–326.
- Evangelista,E.A., Lemaitre,R.N., Sotoodehnia,N., Gharib,S.A. and Totah,R.A. (2018) CYP2J2 expression in adult ventricular myocytes protects against reactive oxygen species toxicity. *Drug Metab. Dispos.*, 46, 380–386.
- 48. Stavropoulou, E., Pircalabioru, G.G. and Bezirtzoglou, E. (2018) The role of cytochromes P450 in infection. *Front. Immunol.*, **9**, 89.
- Liu,Z., Megaraj,V., Li,L., Sell,S., Hu,J. and Ding,X. (2015) Suppression of pulmonary CYP2A13 expression by carcinogen-induced lung tumorigenesis in a CYP2A13-humanized mouse model. *Drug Metab. Dispos.*, 43, 698–702.
- Mazerska,Z., Mroz,A., Pawlowska,M. and Augustin,E. (2016) The role of glucuronidation in drug resistance. *Pharmacol. Ther.*, **159**, 35–55.
- Grayson, P.C., Eddy, S., Taroni, J.N., Lightfoot, Y.L., Mariani, L., Parikh, H., Lindenmeyer, M.T., Ju, W., Greene, C.S., Godfrey, B. *et al.* (2018) Metabolic pathways and immunometabolism in rare kidney diseases. *Ann. Rheum. Dis.*, **77**, 1226–1233.
- Li,Y.H., Li,X.X., Hong,J.J., Wang,Y.X., Fu,J.B., Yang,H., Yu,C.Y., Li,F.C., Hu,J., Xue,W.W. *et al.* (2020) Clinical trials, progression-speed differentiating features and swiftness rule of the innovative targets of first-in-class drugs. *Brief. Bioinform.*, 21, 649–662.
- Barrett, T., Wilhite, S.E., Ledoux, P., Evangelista, C., Kim, I.F., Tomashevsky, M., Marshall, K.A., Phillippy, K.H., Sherman, P.M., Holko, M. *et al.* (2013) NCBI GEO: archive for functional genomics data sets-update. *Nucleic Acids Res.*, 41, D991–D995.
- Gautier, L., Cope, L., Bolstad, B.M. and Irizarry, R.A. (2004) Affy-analysis of affymetrix genechip data at the probe level. *Bioinformatics*, 20, 307–315.
- 55. Bolstad, B.M., Irizarry, R.A., Astrand, M. and Speed, T.P. (2003) A comparison of normalization methods for high density

oligonucleotide array data based on variance and bias. *Bioinformatics*, **19**, 185–193.

- Ge, X., Yamamoto, S., Tsutsumi, S., Midorikawa, Y., Ihara, S., Wang, S.M. and Aburatani, H. (2005) Interpreting expression profiles of cancers by genome-wide survey of breadth of expression in normal tissues. *Genomics*, 86, 127–141.
- Alexander, J.L., Wilson, I.D., Teare, J., Marchesi, J.R., Nicholson, J.K. and Kinross, J.M. (2017) Gut microbiota modulation of chemotherapy efficacy and toxicity. *Nat. Rev. Gastroenterol. Hepatol.*, 14, 356–365.
- Yang,M., Ma,J., Ruan,J., Ye,Y., Fu,P.P. and Lin,G. (2019) Intestinal and hepatic biotransformation of pyrrolizidine alkaloid N-oxides to toxic pyrrolizidine alkaloids. *Arch. Toxicol.*, 93, 2197–2209.
- Dieterich, W., Schink, M. and Zopf, Y. (2018) Microbiota in the gastrointestinal tract. *Med. Sci.*, 6, 116.
- 60. Tang, J., Fu, J., Wang, Y., Li, B., Li, Y., Yang, Q., Cui, X., Hong, J., Li, X., Chen, Y. *et al.* (2020) ANPELA: analysis and performance assessment of the label-free quantification workflow for metaproteomic studies. *Brief. Bioinform.*, 21, 621–636.
- Klatt,N.R., Cheu,R., Birse,K., Zevin,A.S., Perner,M., Noel-Romas,L., Grobler,A., Westmacott,G., Xie,I.Y., Butler,J. *et al.* (2017) Vaginal bacteria modify HIV tenofovir microbicide efficacy in African women. *Science*, **356**, 938–945.
- 62. Tang, J., Fu, J., Wang, Y., Luo, Y., Yang, Q., Li, B., Tu, G., Hong, J., Cui, X., Chen, Y. *et al.* (2019) Simultaneous improvement in the precision, accuracy, and robustness of label-free proteome quantification by optimizing data manipulation chains. *Mol. Cell Proteomics*, 18, 1683–1699.
- Ballesta, A., Innominato, P.F., Dallmann, R., Rand, D.A. and Levi, F.A. (2017) Systems chronotherapeutics. *Pharmacol. Rev*, 69, 161–199.
- 64. Sofi,H.S., Abdal-Hay,A., Ivanovski,S., Zhang,Y.S. and Sheikh,F.A. (2020) Electrospun nanofibers for the delivery of active drugs through nasal, oral and vaginal mucosa: current status and future perspectives. *Mater. Sci. Eng. C Mater. Biol. Appl.*, **111**, 110756.
- Federhen, S. (2015) Type material in the NCBI taxonomy database. Nucleic Acids Res., 43, D1086–D1098.
- 66. Ding,L., Li,L., Liu,S., Bao,X., Dickman,K.G., Sell,S.S., Mei,C., Zhang,Q.Y., Gu,J. and Ding,X. (2020) Proximal tubular vacuolization and hypersensitivity to drug-induced nephrotoxicity in male mice with decreased expression of the NADPH-cytochrome P450 reductase. *Toxicol. Sci.*, **173**, 362–372.
- 67. Yang,Z.Z., Li,L., Wang,L., Yuan,L.M., Xu,M.C., Gu,J.K., Jiang,H.D., Yu,L.S. and Zeng,S. (2017) The regioselective glucuronidation of morphine by dimerized human UGT2B7, 1A1, 1A9 and their allelic variants. *Acta Pharmacol. Sin.*, **38**, 1184–1194.
- Liu, Y.Q., Yuan, L.M., Gao, Z.Z., Xiao, Y.S., Sun, H.Y., Yu, L.S. and Zeng, S. (2016) Dimerization of human uridine diphosphate glucuronosyltransferase allozymes 1A1 and 1A9 alters their quercetin glucuronidation activities. *Sci. Rep.*, 6, 23763.
- 69. Yuan,L.M., Gao,Z.Z., Sun,H.Y., Qian,S.N., Xiao,Y.S., Sun,L.L. and Zeng,S. (2016) Inter-isoform hetero-dimerization of human UDP-glucuronosyltransferases (UGTs) 1A1, 1A9, and 2B7 and impacts on glucuronidation activity. *Sci. Rep.*, 6, 34450.
- Chen,L., Bao,Y., Piekos,S.C., Zhu,K., Zhang,L. and Zhong,X.B. (2018) A transcriptional regulatory network containing nuclear receptors and long noncoding RNAs controls basal and drug-induced expression of cytochrome P450s in HepaRG cells. *Mol. Pharmacol.*, **94**, 749–759.
- Pan,X., Ning,M. and Jeong,H. (2017) Transcriptional regulation of CYP2D6 expression. *Drug Metab. Dispos.*, 45, 42–48.
- Peng,L. and Zhong,X. (2015) Epigenetic regulation of drug metabolism and transport. *Acta Pharm. Sin. B*, 5, 106–112.
- Fisel, P., Schaeffeler, E. and Schwab, M. (2016) DNA methylation of ADME genes. *Clin. Pharmacol. Ther.*, 99, 512–527.
- 74. Mallick, P., Taneja, G., Moorthy, B. and Ghose, R. (2017) Regulation of drug-metabolizing enzymes in infectious and inflammatory disease: implications for biologics-small molecule drug interactions. *Expert Opin. Drug Metab. Toxicol.*, **13**, 605–616.
- 75. Wang,Z., Wu,X. and Wang,Y. (2018) A framework for analyzing DNA methylation data from Illumina Infinium HumanMethylation450 BeadChip. *BMC Bioinformatics*, **19**, 115.
- 76. Jiao, C., Zhang, C., Dai, R., Xia, Y., Wang, K., Giase, G., Chen, C. and Liu, C. (2018) Positional effects revealed in Illumina methylation array and the impact on analysis. *Epigenomics*, 10, 643–659.

- 77. Li,F., Zhou,Y., Zhang,X., Tang,J., Yang,Q., Zhang,Y., Luo,Y., Hu,J., Xue,W., Qiu,Y. *et al.* (2020) SSizer: determining the sample sufficiency for comparative biological study. *J. Mol. Biol.*, **432**, 3411–3421.
- Fu,J., Tang,J., Wang,Y., Cui,X., Yang,Q., Hong,J., Li,X., Li,S., Chen,Y., Xue,W. *et al.* (2018) Discovery of the consistently well-performed analysis chain for SWATH-MS based pharmacoproteomic quantification. *Front. Pharmacol.*, 9, 681.
- Warden, C.D., Lee, H., Tompkins, J.D., Li, X., Wang, C., Riggs, A.D., Yu, H., Jove, R. and Yuan, Y.C. (2013) COHCAP: an integrative genomic pipeline for single-nucleotide resolution DNA methylation analysis. *Nucleic Acids Res.*, 41, e117.
- Song,M.A., Tiirikainen,M., Kwee,S., Okimoto,G., Yu,H. and Wong,L.L. (2013) Elucidating the landscape of aberrant DNA methylation in hepatocellular carcinoma. *PLoS One*, 8, e55761.
- 81. Lancet, T. (2019) ICD-11. Lancet, 393, 2275.
- Piekos, S.C., Chen, L., Wang, P., Shi, J., Yaqoob, S., Zhu, H.J., Ma, X. and Zhong, X.B. (2018) Consequences of phenytoin exposure on hepatic cytochrome P450 expression during postnatal liver maturation in mice. *Drug Metab. Dispos.*, 46, 1241–1250.
- Li,L., Bao,X., Zhang,Q.Y., Negishi,M. and Ding,X. (2017) Role of CYP2B in phenobarbital-induced hepatocyte proliferation in mice. *Drug Metab. Dispos.*, 45, 977–981.
- Backman, J.T., Filppula, A.M., Niemi, M. and Neuvonen, P.J. (2016) Role of cytochrome P450 2C8 in drug metabolism and interactions. *Pharmacol. Rev.*, 68, 168–241.
- Yang,Z., Li,L., Hu,H., Xu,M., Gu,J., Wang,Z.J., Yu,L. and Zeng,S. (2016) Reverse of acute and chronic morphine tolerance by lithocholic acid via down-regulating UGT2B7. *Front. Pharmacol.*, 7, 404.
- 86. Yu, A.M., Ingelman-Sundberg, M., Cherrington, N.J., Aleksunes, L.M., Zanger, U.M., Xie, W., Jeong, H., Morgan, E.T., Turnbaugh, P.J., Klaassen, C.D. *et al.* (2017) Regulation of drug metabolism and toxicity by multiple factors of genetics, epigenetics, lncRNAs, gut microbiota, and diseases: a meeting report of the 21(st) international symposium on microsomes and drug oxidations (MDO). *Acta Pharm. Sin. B*, **7**, 241–248.
- Sun,M., Zhang,Q., Yang,X., Qian,S.Y. and Guo,B. (2016) Vitamin D enhances the efficacy of irinotecan through miR-627-mediated inhibition of intratumoral drug metabolism. *Mol. Cancer Ther.*, 15, 2086–2095.

- Yin,J., Sun,W., Li,F., Hong,J., Li,X., Zhou,Y., Lu,Y., Liu,M., Zhang,X., Chen,N. *et al.* (2020) VARIDT 1.0: variability of drug transporter database. *Nucleic Acids Res.*, 48, D1042–D1050.
- Li,X., Tian,Y., Tu,M.J., Ho,P.Y., Batra,N. and Yu,A.M. (2019) Bioengineered miR-27b-3p and miR-328-3p modulate drug metabolism and disposition via the regulation of target ADME gene expression. *Acta Pharm. Sin. B*, 9, 639–647.
- Li,F., Fan,C., Marquez-Lago,T.T., Leier,A., Revote,J., Jia,C., Zhu,Y., Smith,A.I., Webb,G.I., Liu,Q. *et al.* (2020) PRISMOID: a comprehensive 3D structure database for post-translational modifications and mutations with functional impact. *Brie.f Bioinform.*, 21, 1069–1079.
- Yang, Q., Wang, Y., Zhang, Y., Li, F., Xia, W., Zhou, Y., Qiu, Y., Li, H. and Zhu, F. (2020) NOREVA: enhanced normalization and evaluation of time-course and multi-class metabolomic data. *Nucleic Acids Res.*, 48, W436–W448.
- 92. Xue, W., Yang, F., Wang, P., Zheng, G., Chen, Y., Yao, X. and Zhu, F. (2018) What contributes to serotonin-norepinephrine reuptake inhibitors' dual-targeting mechanism? The key role of transmembrane domain 6 in human serotonin and norepinephrine transporters revealed by molecular dynamics simulation. ACS Chem. Neurosci., 9, 1128–1140.
- Li,B., Tang,J., Yang,Q., Li,S., Cui,X., Li,Y., Chen,Y., Xue,W., Li,X. and Zhu,F. (2017) NOREVA: normalization and evaluation of MS-based metabolomics data. *Nucleic Acids Res.*, 45, W162–W170.
- 94. Yang, Q., Li, B., Tang, J., Cui, X., Wang, Y., Li, X., Hu, J., Chen, Y., Xue, W., Lou, Y. *et al.* (2020) Consistent gene signature of schizophrenia identified by a novel feature selection strategy from comprehensive sets of transcriptomic data. *Brief. Bioinform.*, 21, 1058–1068.
- Ning, B., Yu, D. and Yu, A.M. (2019) Advances and challenges in studying noncoding RNA regulation of drug metabolism and development of RNA therapeutics. *Biochem Pharmacol*, 169, 113638.
- Zhang, Y., Xie, R., Wang, J., Leier, A., Marquez-Lago, T.T., Akutsu, T., Webb, G.I., Chou, K.C. and Song, J. (2019) Computational analysis and prediction of lysine malonylation sites by exploiting informative features in an integrative machine-learning framework. *Brief. Bioinform.*, 20, 2185–2199.